# Landbird Monitoring in Denali National Park and Preserve

# Objective

Determine long-term trends in avian abundance.

## **Design Considerations**

- Sampling methodology
- Independence of sampling units
- Observer bias
- Sample size and power to detect trends
- Geographic scale

## **Strengths of Protocol**

- 90% chance of detecting a 50% decline in a 25-year period.
- 12 species of forest songbirds.
- Robust design tailored to Denali.
- In line with the LTEM management focus objective.

### Concerns

Are 12 species enough?

• Spatial scale of sampling is not representative of Denali Park.


Debevec/Rexstad analysis.

# **Ecological Focus Objective**

To improve the understanding of the Denali ecosystem.

# Could songbird declines affect ecosystem function in Denali Park?

# What role do forest birds play in ecosystem function?



#### **Nesting Success**

**Predators** Weather

**Insect Abundance** 

**Plant Biomass** 

\*Test assumptions of abundance estimates

### Recommendations

### **Management Focus Objective**

- Conduct surveys in habitats representative of Denali Park.
- Evaluate whether more species should be monitored.
- Determine whether assumptions of abundance trends are correct.

### Recommendations

### **Ecological Focus Objective**

• Examine factors affecting nest success.

 Study the role songbirds play in ecosystem processes in Denali.